AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:

LISTING OF CLAIMS:

1. (Withdrawn) A method for treating diseases caused by and/or associated with an altered protein kinase activity which comprises administering to a mammal in need thereof an effective amount of a pyrrolo-pyrazole or pyrazolo-azepine derivative represented by formula (I):

$$\begin{array}{c|c} R & R_2 \\ \hline (CH_2)_m & (CH_2)_n \\ \hline R_d & R_b \\ \hline R_c & R_a \\ \end{array}$$
 (I)

wherein R represents hydrogen or halogen atom, or an optionally substituted group selected from aryl C₂-C₆ alkenyl, (heterocyclyl) C₂-C₆ alkenyl, aryl C₂-C₆ alkynyl, or (heterocyclyl) C₂-C₆ alkynyl group, -R', -COR', -COOR', -CN, -CONR'R", -OR', -S(O)_qR', -SO₂NR'R", -B(OR"')₂, -SnR"", wherein R' and R", the same or different, independently represent hydrogen atom or an optionally further substituted straight or branched C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, saturated or unsaturated C₃-C₆ cycloalkyl, aryl, heterocyclyl, aryl C₁-C₆ alkyl or (heterocyclyl)C₁-C₆ alkyl; R" represents hydrogen, C₁-C₆ alkyl, or R"', together with the two oxygen and the boron atoms, forms a saturated or unsaturated C₅-C₈ (hetero)cycloalkyl, optionally benzocondensed or substituted, and R"" represents C₁-C₆ alkyl;

R₁ represents hydrogen atom or an optionally substituted group selected from -R',

-CH₂R', -COR', -COOR', -CONR'R", -C(=NH)NHR', -S(O)_qR', or -SO₂NR'R", wherein R' and R" are as defined above;

 R_2 represents hydrogen atom, -COR', -COOR', -CONR'R", -S(O)_q R', -SO₂NR'R", C₁-C₆ alkyl or (heterocyclyl)C₁-C₆ alkyl group, wherein R' and R" are as defined above;

 R_a , R_b , R_c and R_d , being the same or different, independently represent hydrogen atom, an optionally further substituted straight or branched C_1 - C_6 alkyl, aryl, heterocyclyl, aryl C_1 - C_6 alkyl, (heterocyclyl) C_1 - C_6 alkyl or - CH_2OR ' group, wherein R' is as above defined, or R_a and R_b and/or R_c and R_d , taken together with the carbon atom to which they are bonded, form an optionally substituted, saturated or unsaturated, C_3 - C_6 cycloalkyl group; q is 0, 1 or 2; m and n, each independently, represents 0, 1 or 2, provided that m + n is 0 or equal to 2; or a pharmaceutically acceptable salt thereof.

- 2. (Withdrawn) The method of claim 1 wherein the disease caused by and/or associated with an altered protein kinase activity is selected from the group consisting of cancer, cell proliferative disorders, Alzheimer's disease, viral infections, auto-immune diseases and neurodegenerative disorders.
- 3. (Withdrawn) The method of claim 2 wherein the cancer is selected from carcinoma, squamous cell carcinoma, hematopoietic tumors of myeloid or lymphoid lineage, tumors of mesenchymal origin, tumors of the central and peripheral nervous system, melanoma, seminoma, teratocarcinoma, osteosarcoma, xeroderma pigmentosum, keratocanthoma, thyroid follicular cancer and Kaposi's sarcoma.

- 4. (Withdrawn) The method of claim 2 wherein the cell proliferative disorder is selected from the group consisting of benign prostate hyperplasia, familial adenomatosis polyposis, neuro-fibromatosis, psoriasis, vascular smooth cell proliferation associated with atherosclerosis, pulmonary fibrosis, arthritis glomerulonephritis and post-surgical stenosis and restenosis.
- 5. (Withdrawn) The method of claim 1 which provides tumor angiogenesis and metastasis inhibition.
- 6. (Withdrawn) The method of claim 1 further comprising subjecting the mammal in need thereof to a radiation therapy or chemotherapy regimen in combination with at least one cytostatic or cytotoxic agent.
 - 7. (Withdrawn) The method of claim 1 wherein the mammal in need thereof is a human.
- 8. (Withdrawn) The method of claim 1 wherein in the compound of formula (I) R is H, I, Br, Cl, F, aryl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, -B(OR"')₂, -COR', -CONR'R", -CN, SO₂R', OR', SR', and R₁ is H, C₁-C₆ alkyl, aryl, -COR', -CONR'R", -COOR', -SO₂R', or -SO₂NR'R", and R₂ is H, -COOR', -COR', -CONR'R", C₁-C₆ alkyl, -SO₂R', or -SO₂NR'R", (heterocyclyl) C₁-C₆ alkyl group , wherein R' and R", the same or different, are selected from hydrogen or optionally substituted straight or branched C₁-C₆ alkyl, aryl or aryl C₁-C₆ alkyl groups;

 R_a , R_b , R_c and R_d , the same or different, are selected from hydrogen or straight or branched C_1 - C_3 alkyl or, taken together with the carbon atom to which they are bonded form a C_3 - C_6 cycloalkyl group.

- 9. (Withdrawn) The method of claim 1 wherein, in the compound of formula (I), R is selected from aryl, -COR', -CONR'R", wherein R' and R", the same or different, are selected from hydrogen or optionally substituted straight or branched C₁-C₆ alkyl, aryl or aryl C₁-C₆ alkyl groups.
- 10. (Withdrawn) The method of claim 1 wherein, in the compound of formula (I), R₁ is selected from H, C₁-C₆ alkyl, aryl, -COR', -CONR'R", COOR', -SO₂R' or -SO₂NR'R", wherein R' and R", the same or different, are selected from hydrogen or optionally substituted straight or branched C₁-C₆ alkyl, aryl or aryl C₁-C₆ alkyl groups.
- 11. (Withdrawn) The method of claim 1 wherein, in the compound of formula (I), R₂ is H,
 -COOR', -CONR'R", C₁-C₆ alkyl, wherein R' and R", the same or different, are selected from hydrogen or optionally substituted straight or branched C₁-C₆ alkyl, aryl or aryl C₁-C₆ alkyl groups.
- 12. (Withdrawn) A method for inhibiting protein kinase activity which comprises contacting the said kinase with an effective amount of a compound of formula (I) as defined in claim 1.

13. (Currently Amended) A pyrrolo-pyrazole or pyrazolo-azepine derivative represented by formula (I):

wherein R is aryl

 R_1 represents hydrogen atom or an optionally substituted group selected from -R', $-CH_2R'$, -COR', -COOR', -CONR'R'', -C(=NH)NHR', $-S(O)_qR'$, or $-SO_2NR'R''$, wherein R' and R'', the same or different, independently represent hydrogen atom or an optionally further substituted straight or branched C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, saturated or unsaturated C_3 - C_6 cycloalkyl, aryl, heterocyclyl, aryl C_1 - C_6 alkyl or (heterocyclyl) C_1 - C_6 alkyl with the proviso that when R_1 is -R', R' is not heterocyclyl or (heterocyclyl) C_1 - C_6 alkyl;

R₂ represents hydrogen atom, -COR', -COOR', -CONR'R", -S(O)_q R', -SO₂NR'R", or C₁-C₆ alkyl, or (heterocyclyl)C₁-C₆ alkyl group, wherein R' and R" are as defined above;

 R_a , R_b , R_c and R_d , being the same or different, independently represent hydrogen atom, an optionally further substituted straight or branched C_1 - C_6 alkyl, aryl, aryl C_1 - C_6 alkyl, or - CH_2OR ' group, wherein R' is as above defined, or R_a and R_b and/or R_c and R_d , taken together with the carbon atom to which they are bonded, form an optionally substituted, saturated or unsaturated, C_3 - C_6 cycloalkyl group; q is 0, 1 or 2; m and m, each independently, represents 0, 1 or 2, provided that m+n is 0; and with the further proviso that R_1 and R_2 are not heterocyclylalkyl;

or a pharmaceutically acceptable salt thereof.

14. (Previously Presented) A compound of formula (I) according to claim 13 wherein R₁ is H, C₁-C₆ alkyl, aryl, -COR', -CONR'R", -COOR', -SO₂R', or -SO₂NR'R", and R₂ is H, -COOR', -CONR'R", C₁-C₆ alkyl, -SO₂R', or -SO₂NR'R", wherein R' and R", the same or different, are selected from hydrogen or optionally substituted straight or branched C₁-C₆ alkyl, aryl or aryl C₁-C₆ alkyl groups;

 R_a , R_b , R_c and R_d , the same or different, are selected from hydrogen or straight or branched C_1 - C_3 alkyl or, taken together with the carbon atom to which they are bonded form a C_3 - C_6 cycloalkyl group.

15. (Previously Presented) A compound of formula (I) according to claim 13 wherein R' and R", the same or different, are selected from hydrogen or optionally substituted straight or branched C_I-C₆ alkyl, aryl or aryl C₁-C₆ alkyl groups.

16. (Cancelled)

- 17. (Original) A compound of formula (I) according to claim 13 wherein R₂ is H, -COOR', -CONR'R", C₁-C₆ alkyl, wherein R' and R", the same or different, are selected from hydrogen or optionally substituted straight or branched C₁-C₆ alkyl, aryl or aryl C₁-C₆ alkyl groups.
- 18. (Withdrawn) A process for preparing the compounds of formula (I) or the pharmaceutically acceptable salts thereof, as defined in claim 13, which process comprises:
 - a) submitting a compound of formula (II)

$$H_2N$$
 N
 R_2
 $(CH_2)_m$
 $(CH_2)_n$
 R_b
 R_1
 (II)

wherein R₁ is as defined in claim 13 but not hydrogen atom, and R_a, R_b, R_c, R_d, R₂, m and n are as defined in claim 13, to diazotation and subsequent appropriate quenching, thus obtaining a compound of formula (I)

wherein R₁ is as defined above but not hydrogen; R_a, R_b, R_c, R_d, R₂, m and n are as defined above, and R is hydrogen, iodine, bromine, chlorine or fluorine atom or a CN group;

b1) converting a thus obtained compound of formula (I) wherein R is I, Br, Cl into another compound of formula (I) wherein R is an optionally substituted aryl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, -SR', -OR' or -COR' wherein R' is as defined in claim 13;

b2) converting a compound of formula (I) wherein R is hydrogen into another compound of formula (I) wherein R is -B(OR'")₂, -SnR'", -COOR', -COR', C₁-C₆ alkyl or iodine, wherein R', R'" and R'" are as defined in claim 13;

c) converting a compound of formula (I) wherein R is -B(OR'")₂ or -SnR'" as above defined into another compound of formula (I) wherein R is an optionally substituted aryl, C₂-C₆ alkynyl;

d) optionally converting a compound of formula (I) into another different compound of formula (I), and, if desired, converting a compound of formula (I) into a pharmaceutically acceptable salt thereof or converting a salt into the free compound (I).

19. (Withdrawn) A process for preparing a compound of formula (I) according to claim 13, which which process comprises:

either b1a) converting a compound of formula (I) into another compound of formula (I) wherein R has the meanings of claim 18 resulting from step b1 and R₁, R_a, R_b, R_c, R_d, m and n are as defined in claim 13, analogously to step b1 defined in claim 18 and Pa) reacting the resultant compound of formula (I) wherein R, R_a, R_b, R_c, R_d, m and n are as defined above, R₁ is as described above but not hydrogen and R₂ is hydrogen, with a suitable solid support so as to obtain a compound of formula (III)

wherein R, R_a , R_b , R_c , R_d , m and n are as defined above, R_1 is as defined in claim 13 but not hydrogen, and Q is a solid support, or P) reacting a compound of formula (I) wherein R, R_a , R_b , R_c , R_d , m and n are as defined above, R_1 is as defined above but not hydrogen and R_2 is

hydrogen, with a suitable solid support so as to obtain a compound of formula (III) as defined above and

- B) then, analogously to steps b1, b2, c and d defined in claim 18, optionally converting a thus obtained compound of formula (III) into another compound of formula (III) wherein R has the meanings defined in claim 18 for steps b1 to d and R₁, R_a, R_b, R_c, R_d, m and n are as defined above;
- C) cleaving a compound of formula (III) so as to eliminate the solid support and to obtain the desired compound of formula (I);
- D) optionally converting a compound of formula (I) into another different compound of formula (I),

and, if desired, converting a compound of formula (I) into a pharmaceutically acceptable salt thereof or converting a salt into the free compound (I) as described above.

20. (Previously Presented) A compound of formula (III)

wherein R₁, R, R_a, R_b, R_c, R_d, m and n are as defined in claim 13, and Q is a solid support wherein said solid support is a residue derived from a resin selected from the group consisting of isocyanate polystyrenic resin, 2-chloro-trityl chloride resin, trityl chloride resin, p-nitrophenyl carbonate Wang resin and the bromo-4-methoxyphenyl)methyl polystyrene.

21. (Cancelled)

22. (Withdrawn) A process for preparing a compound of formula (III) as defined in claim 20, which process comprises:

either b1a) converting a compound of formula (I) into another compound of formula (I) wherein R is as defined in claim 19 resulting from step b1 and R₁, R_a, R_b, R_c, R_d, m and n are as defined in claim 13, analogously to step b1 described in claim 18 and Pa) reacting the resultant compound of formula (I) wherein R, R_a, R_b, R_c, R_d, m and n are as defined above, R₁ is as defined above but not hydrogen and R₂ is hydrogen, with a suitable solid support so as to obtain a compound of formula (III)

$$\begin{array}{c|c}
R & & & & & Q \\
\hline
(CH_2)_m & & & & (CH_2)_n \\
R_d & & & & & & R_b \\
R_c & & & & & & R_a
\end{array}$$
(III)

wherein R, R_a , R_b , R_c , R_d , m and n are as defined above, R_1 is as defined in claim 13 but not hydrogen, and Q is a solid support, or

A) reacting a compound of formula (I) wherein R, R_a, R_b, R_c, R_d, m and n are as defined above, R₁ is as defined above but not hydrogen and R₂ is hydrogen, with a suitable solid support so as to obtain a compound of formula (III) as defined above and

B) then, analogously to steps b1, b2, c and d described in claim 18, optionally converting a thus obtained compound of formula (III) into another compound of formula (III) wherein R has

the meanings as defined in claim 18 for steps b1 to d and R₁, R_a, R_b, R_c, R_d, m and n are as defined above.

23. (Original) A library of two or more compounds of formula (I):

wherein R, R_1 , R_2 R_a , R_b , R_c , R_d m and n are as defined in claim 13, which can be obtained starting from one or more compound supported onto a solid support of the formula (III) as defined in claim 20.

- 24. (Cancelled)
- 25. (Original) A pharmaceutical composition comprising a therapeutically effective amount of a compound of formula (I), as defined in claim 13, and at least one pharmaceutically acceptable carrier and/or diluent.
 - 26. (Cancelled)
- 27. (Currently Amended) A product comprising a compound of formula (I) as defined in claim 13 or a pharmaceutical composition thereof as defined in claim 25, and one or more chemotherapeutic agents, as a combined preparation for simultaneous, separate or sequential use in anticancer therapy wherein the cancer is selected from carcinoma, squamous cell carcinoma,

hematopoietic tumors of myeloid or lymphoid lineage, tumors of mesenchymal origin, tumors of the central and peripheral nervous system, melanoma, seminoma, teratocarcinoma, osteosarcoma, xeroderma pigmentosum, keratocanthoma, thyroid follicular cancer and Kaposi's sarcoma.

- 28. (Cancelled)
- 29. (Withdrawn) Use of a compound of formula (I), as defined in claim 1, in the manufacture of a medicament with antitumor activity.